



# MURANG'A UNIVERSITY OF TECHNOLOGY

## SCHOOL OF PURE AND APPLIED SCIENCE

DEPARTMENT OF APPLIED SCIENCE

UNIVERSITY ORDINARY EXAMINATION

2017/2018 ACADEMIC YEAR

**FIRST YEAR SECOND SEMESTER EXAMINATION FOR BACHELOR OF  
SCIENCE IN APPLIED STATISTICS WITH PROGRAMMING**

AMS107: PROBABILITY AND DISTRIBUTION THEORY I

DURATION: 2 HOURS

DATE: 26<sup>TH</sup> APRIL 2018

TIME: 8.00AM – 11.00AM

### **Instructions to Candidates:**

1. Answer **Section A** and **Any Other Two** questions in **Section B**.
2. Mobile phones are not allowed in the examination room.
3. You are not allowed to write on this examination question paper.

## SECTION A – ANSWER ALL QUESTIONS IN THIS SECTION (30 Marks)

### QUESTION ONE

- a) Define the following terms as used in probability and statistics
- i. Random variable (1 Mark)
  - ii. Probability distribution (1 Mark)
  - iii. Random Vector (1 Mark)
  - iv. Correlation coefficient (1 Mark)

- b) If  $x$  is a discrete random variable
- i. From a Bernoulli distribution given as

$$f(x, \theta) = \begin{cases} u^x (1 - u)^{1-x} & x = 0, 1 \\ 0 & \text{otherwise} \end{cases}$$

Derive the moment generating function of  $x$  (4 Marks)

- ii. Given a random variable  $x$  a POPn

$$f(x) = \begin{cases} 1 & \text{for } -1/2 < x < \theta + 1/2 \\ 0 & \text{otherwise} \end{cases}$$

Obtain the expected value of  $x$ . (4 Marks)

- c) i) Define the moment generating function of a continuous random variable  $x$ . (2 Marks)

Hence or otherwise obtain the moment generating function of a random variable

$X \sim N(0, 1)$  (5 Marks)

- ii) Obtain the moment generating function of a random variable  $Y$  which is normal with the mean  $\mu$  and standard deviation  $\delta$  using results of (i) above. (4 Marks)

- d) suppose  $X$  and  $Y$  are two continuous random variable with a bivariate probability density function given by

$$f_{xy}^{(x,y)} = \begin{cases} x = y & \text{if } 0 \leq x, y < 1 \\ 0 & \text{elsewhere} \end{cases}$$

- i. Show that

$$\iint_{IR} f_{xy}^{(x,y)} dx dy = 1$$

(3 Marks)

- ii. Obtain the marginal probability density of  $X$  ( $f_x^{(x)}$ ) (4 Marks)

## SECTION B – ANSWER ANY TWO QUESTIONS IN THIS SECTION

### QUESTION TWO (20 MARKS)

- a) Define the stochastic independence of two random variables (3 Marks)
- b) Let  $X$  and  $Y$  have  $C$  bivariate probability distribution given as

$$f_{xy}^{(x,y)} = \begin{cases} 4xy + 2x + 2y + 1/4 & \text{if } 0 \leq x, y < 1 \\ 0 & \text{Otherwise} \end{cases}$$

- i. Obtain the marginal distribution of  $X$  and  $Y$  (4 Marks)

- ii. Determine whether  $X$  and  $Y$  are stochastically independent. (4 Marks)
- c) Suppose  $X$  and  $Y$  are two independent standard normal random variables, determine the joint probability density of  $X$  and  $Y$  i.e.  $f_{xy}^{(x,y)}$  (7 Marks)

**QUESTION THREE (20 MARKS)**

- a) State any two properties of a gamma distribution. (4 Marks)
- b) A random variable  $X$  has a gamma distribution of its probability distribution given as

$$f(t) = \begin{cases} \frac{\lambda^\alpha}{(\alpha)} t^{\alpha-1} e^{-\lambda t} & \text{for } t \geq 0 \\ 0 & \text{elsewhere} \end{cases} \quad \alpha, \lambda > 0$$

- i. Obtain the expectation and variance of  $X$  by direct method (or using *pdf*)(12 Marks)
- ii. Obtain the moment generating function of  $X$  (4 Marks)

**QUESTION FOUR (20 MARKS)**

- a) Let  $X$  be a random variable with a density function

$$f(x) = \begin{cases} x^3/3 & \text{for } -1 < x < 2 \\ 0 & \text{Otherwise} \end{cases}$$

Find the expected value of  $g(x) = 4x + 3$  (5 Marks)

- b) A continuous random variable  $X$  has a moment generating function given by

$$M_X(t) = \frac{9}{(3-t)^2}$$

Find the mean and variance of  $X$  (6 Marks)

- c) A random variable  $X$  is said to have a chi-square distribution with  $n$ -degrees of freedom if its mgf (moment generating function) is defined for any  $t < 1/2$  and it's equal to  $M_x(t) = (1 - 2t)^{-n/2}$ . If  $Y = X_1 + X_2$  where  $X_1$  and  $X_2$  are independent random variables have chi-square distribution with  $n_1$  and  $n_2$  degrees of freedom respectively. Prove that  $X$  has chi-square with  $n_1 + n_2$  *dof*. (9 Marks)